



ELSEVIER

 JOURNAL OF
 ADOLESCENT
 HEALTH

www.jahonline.org

Review article

A Package of Sexual and Reproductive Health and Rights Interventions—What Does It Mean for Adolescents?



Danielle Marie Claire Engel, M.A.^{a,*}, Mandira Paul, Ph.D.^a, Satvika Chalasani, Ph.D.^a, Lianne Gonsalves, M.S.P.H.^b, David Anthony Ross, M.A., M.Sc., B.M.B.Ch., Ph.D.^c, Venkatraman Chandra-Mouli, M.B.B.S., M.Sc.^b, Claire B. Cole, M.P.H.^d, Cristina de Carvalho Eriksson, Ph.D.^e, Brendan Hayes, M.Sc.^f, Anandita Philipose^g, Sally Beadle, M.P.H.^h, and B. Jane Ferguson, M.S.W., M.Sc.ⁱ

^a Technical Division, United Nations Population Fund, New York, New York

^b Department of Reproductive Health and Research, World Health Organization/Human Reproduction Programme, Geneva, Switzerland

^c Department of Maternal, Newborn, Child, and Adolescent Health, World Health Organization, Geneva, Switzerland

^d Population Services International, Washington, DC

^e Health Section, Programme Division, United Nations Children Fund, New York, New York

^f Global Financing Facility, World Bank, Washington, DC

^g Eastern and Southern Regional Office, United Nations Population Fund, Johannesburg, South Africa

^h Section of Health and Education, United Nations Educational, Scientific and Cultural Organization, Paris, France

ⁱ Independent Consultant, Adolescent Health and Development, Tannay, Switzerland

Article history: Received July 17, 2019; Accepted September 24, 2019

Keywords: Adolescent sexual and reproductive health; Health policy; Comprehensive SRHR package; UHC; Health systems; Delivery models; Adolescent responsive health services; Determinants of ASRH

ABSTRACT

This article analyzes the relevance of the comprehensive definition of sexual and reproductive health and rights (SRHR) to adolescents and identifies adolescent-specific implications for the implementation of an essential package of SRHR interventions. The delivery of a comprehensive approach to SRHR targeting adolescents is underpinned by five principles—equity, quality, accountability, multisectorality, and meaningful engagement. All SRHR interventions included in the package are relevant to adolescents, given the diversity of adolescents' SRHR needs and considering their specific attributes, circumstances, and experiences. Ensuring that this package is available, accessible, and acceptable to adolescents requires an approach that looks at adolescents as being biologically and socially distinct from other age groups and acknowledges that they face some specific barriers when accessing SRHR services. This article provides cross-cutting strategies for the implementation of a comprehensive approach to SRHR for adolescents and specific considerations in delivering each intervention in the package of essential SRHR interventions. To further implement the International Conference on Population and Development Programme of Action, a prerequisite for achieving the Sustainable Development Goals, SRHR interventions must be adolescent responsive, delivered through multiple

IMPLICATIONS AND CONTRIBUTION

This article presents adolescent-specific considerations for implementing a package of essential SRHR interventions. Adolescents' unique attributes and needs, and the barriers they face in accessing care impact the delivery of each intervention of the package. While certain aspects of SRHR are unique to this age group,

Conflicts of Interest: The authors declare that there are no conflicts of interest.

Disclaimer: Publication of this article was supported by the World Health Organization and the United Nations Population Fund. The opinions or views expressed in this article are those of the authors and do not necessarily represent the official position of the funders.

* Address correspondence to: Danielle Marie Claire Engel, M.A., United Nations Population Fund, 605 Third Avenue, New York, New York, 10158.

E-mail address: engel@unfpa.org (D.M.C. Engel).

platforms, leveraging multisectoral collaboration, and strengthening accountability and participation.

© 2019 Published by Elsevier Inc. on behalf of Society for Adolescent Health and Medicine.

all SRHR interventions are relevant to adolescents and can be adapted to ensure service delivery is adolescent responsive in practice, and contributing to the progressive realization of UHC.

A breakthrough at the 1994 International Conference on Population and Development (ICPD) was the recognition that sexual and reproductive health (SRH) services are essential for all people, including adolescents and youth (This paper defines adolescents as those aged 10–19 years, youth as 15–24 years, and young people as 10–24 years). The ICPD Programme of Action, was forward-looking in relation to adolescents and young people, it called for “meeting the educational and (health) service needs of adolescents to enable them to deal in a positive and responsible way with their sexuality” [1].

In the decades since, the urgency to address the sexual and reproductive health and rights (SRHR) of adolescents and young people has been repeatedly reaffirmed [2,3]. Buttressed by increased knowledge about adolescence development [3,4], there is now near-universal consensus that promoting and protecting adolescent health will bring health benefits for the adolescent now, in their future lives and for the future

generations [3]. Still, ensuring the SRHR of adolescents remains an unfinished agenda in the ICPD Programme of Action [5].

The increased global attention and accumulating evidence related to adolescent SRHR are seldom reflected in robust, scaled-up programmatic responses at the service delivery level [6]. Adolescents risk falling into a policy and service delivery gap, where their specific health needs and barriers to access are overlooked [7].

To respond effectively and adequately to adolescents' health needs, an integrated approach to SRHR (Box 1), implemented through a package of essential SRHR interventions, is needed [8]. Such a package includes promotive, preventive, and curative interventions, taking the social determinants of health into account in its implementation.

Contribution of this Article

This article presents adolescent-specific considerations for implementing the package of essential SRHR interventions as identified in the World Health Organization reproductive health framework [9] and the Guttmacher-Lancet Commission's definition of SRHR [8]. It outlines adolescents' unique attributes and needs and the barriers they face and goes on to identify corresponding strategies and approaches to address these. It aims to guide countries seeking to enhance investments in SRHR tailored to adolescents and progressively realize Universal Health Coverage (UHC) by prioritizing, costing, financing, and implementing the package of essential SRHR interventions for adolescents in all their diversity.

Principles underpinning the package of essential SRHR interventions for adolescents

A human rights-based approach to adolescent health is essential to ensure that adolescents can exercise their rights and make informed choices in relation to their sexual and reproductive health. It obliges duty bearers to respect, protect, and fulfill human rights by refraining from denying or limiting access to health care services or withholding information (including parental consent requirements), taking the necessary legislative and other measures to protect adolescents from discrimination, gender-based violence and harmful practices and to adopt/adapt appropriate legislative, administrative, budgetary, judicial, promotional, and other measures to fully realize adolescents' right to health including sexual and reproductive health [10].

The ICPD Programme of Action highlighted the principles of equity, quality, and accountability as essential for the delivery of SRH services and the realization of individuals' sexual and reproductive rights [1]. Since the ICPD, the importance of multisectorality and meaningful participation of rights holders has been increasingly acknowledged [11]. This article thus outlines

Box 1. Guttmacher–Lancet Commission, The Lancet, 2018. Integrated definition of sexual and reproductive health and rights

Sexual and reproductive health is a state of physical, emotional, mental, and social well-being in relation to all aspects of sexuality and reproduction, not merely the absence of disease, dysfunction, or infirmity. Therefore, a positive approach to sexuality and reproduction should recognize the part played by pleasurable sexual relationships, trust, and communication in the promotion of self-esteem and overall well-being.

All individuals have a right to make decisions governing their bodies and to access services that support that right. Achievement of sexual and reproductive health relies on the realization of sexual and reproductive rights, which are based on the human rights of all individuals to:

- have their bodily integrity, privacy, and personal autonomy respected;
- freely define their own sexuality, including sexual orientation and gender identity and expression;
- decide whether and when to be sexually active;
- choose their sexual partners;
- have safe and pleasurable sexual experiences;
- decide whether, when, and whom to marry;
- decide whether, when, and by what means to have a child or children, and how many children to have;
- have access over their lifetimes to the information, resources, services, and support necessary to achieve all the above, free from discrimination, coercion, exploitation, and violence.

Box 2. Diversity of adolescents and their sexual and reproductive health and rights (SRHR) needs	
Ecological level	Examples of adolescents' circumstances and experiences influencing their SRHR needs
Individual	<p><i>Adolescents who are:</i></p> <ul style="list-style-type: none"> - Very young (aged 10–14 years) - Unmarried and sexually active - Pregnant - First-time young parents - Suffering obstetric fistula - Lesbian, gay, bisexual, transgender, intersex and other sexuality, sex, and gender-diverse adolescents - Living with HIV and/or other chronic illnesses or disabilities; - In need of mental health and psychosocial support services - Out of school or with low level of educational attainment
Interpersonal	<p><i>Adolescents who are:</i></p> <ul style="list-style-type: none"> - Acting as head of households - Married in childhood - Orphans living in host families - Living in households at or below the poverty line - Experiencing or witnessing interpersonal and/or sexual and gender-based violence - victims of trafficking and sexual abuse and exploitation
Community	<p><i>Adolescents living in:</i></p> <ul style="list-style-type: none"> - Urban slum communities, rural remote, and/or pastoralist communities - Camps or settlements as a result of crises
Macro	<p><i>Adolescents in societies affected by:</i></p> <ul style="list-style-type: none"> - Conflict - Disaster - Climate crises - Forced migration <p><i>Adolescents who are:</i></p> <ul style="list-style-type: none"> - Members of ethnic minorities - Refugees - Internally displaced

five key principles that must be considered in the delivery of a package of essential SRHR interventions focusing on how they relate specifically to adolescents.

- Equity in access to SRHR for adolescents means that every adolescent has access to high-quality information and the full range of services to cater to their SRHR needs in a timely manner. It ensures that interventions are inclusive of those marginalized and considers the diversity of adolescents (Box 2). This requires using various delivery platforms to ensure inclusivity, that is, going beyond health facilities and schools, expanding community-based service delivery, including mobile and household-level outreach, as well as reducing or eliminating any financial or physical barriers [12].

- Quality of care has implications across the entire health system [13,14]. Although important for everyone, quality of care has particular implications for service delivery to adolescents. Services must be respectful, nonjudgmental, confidential, and present information that is comprehensible to adolescents to allow for informed decision-making [15]. This requires that providers are trained to provide accurate and nonjudgmental SRHR services and information tailored to adolescents' developmental stage. Variability in quality can be minimized by adopting global standards for quality health care services for adolescents [13] and supporting their achievement by addressing inability to physically access services, out-of-pocket payments, unregulated private health care provision, lack of service demand, and availability of data pertaining to adolescents [6].
- Accountability: A key component of ensuring quality and stigma-free services is to create platforms for feedback, where clients can hold health care providers and decision-makers accountable. Allowing adolescents to engage in accountability mechanisms can be a constructive way to monitor the quality of service provision, identify gaps and traits particularly appreciated by adolescents in different contexts. Such mechanisms can include community scorecard approaches [16], community-facility health management committees, and facility boards that ensure youth representation [3]. Moreover, health management information systems must collect age and gender-disaggregated data on adolescents, including very young adolescents (aged 10–14 years), to allow effective planning, monitoring and evaluation of service delivery. Progress or lack thereof should be documented and transparently shared with communities including adolescents to allow for informed feedback and continuous improvement.
- Multisectorality recognizes that health outcomes are determined by factors both within and outside the health sector and that some interventions critical to the health of adolescents are delivered by other sectors [17]. A multisectoral approach requires programs to build synergies across sectors, such as education, health, labor, transportation, infrastructure, and the social services [11,18–20] and, where relevant, establish coordination mechanisms between line ministries. The Sustainable Development Goals provide an excellent opportunity for tracking multisectoral action, identifying the linkages between the sectors relevant to adolescent SRHR [21].
- Meaningful youth engagement: Adolescents and youth have the right to actively and meaningfully participate in all matters that affect their lives [22]. In addition, greater engagement of adolescents in their health is positively associated with utilization of care and can enhance the effectiveness and sustainability of adolescent SRHR interventions [23–25]. Policy-makers and health service planners need to go beyond the recognition and identification of young people solely as beneficiaries and toward engaging them as equal and valuable partners in decision-making with other stakeholders [26]. This must go hand in hand with positive youth development approaches [27].

Unpacking the Heterogeneity of Adolescents' and Their SRHR Needs

Adolescent behavior and development are shaped by the immediate and distal environmental contexts in which they live

Table 1

Adolescent specific considerations for the delivery of the package of essential sexual and reproductive health and rights (SRHR) interventions

SRHR intervention	Related adolescent-specific attributes, needs, and barriers	Implication for service delivery
Comprehensive Sexuality Education (in and out of school)	Many adolescents lack basic knowledge of SRHR [30–32].	A sexuality education curriculum that is comprehensive, scientifically accurate, timely and developmentally appropriate, and human rights based should be delivered to all adolescents [33].
	In many countries, a significant proportion of adolescents are out of school [34].	The education sector plays a critical role in the provision of CSE, including building the capacity of teachers to deliver effective programs [33].
	Adolescents receive no or very little information from their parents and caregivers [35].	CSE programmes should be introduced in nonformal education and community-based settings to reach out-of-school adolescents including those most vulnerable and marginalized, especially in countries where school attendance is low or where adequate CSE is not included as part of the national curriculum [33].
	Adolescent girls are often uninformed and unprepared for menarche [37].	Forging a common understanding of CSE, building broad social and political consensus around it, and generating support among stakeholders, including opposition groups, requires an ongoing and adequately funded advocacy strategy targeting teachers, parents, religious and community leaders, and policy makers [36].
	Most young people around the world become sexually active during adolescence—whether within marriage or before and whether consensual or forced [38].	CSE can teach adolescents about the stages of the menstrual cycle, including the time around which pregnancy is most able to occur, how to manage menstruation, and that it is a normal and natural part of a girls' physical development and should not be treated with secrecy or stigma [33].
	Adolescence is a time of continued gender socialization. Evidence links traditional gender norms unequal power in sexual relationships and intimate partner violence with negative SRH outcomes [42,43].	CSE needs to start at an early age (ideally before the onset of adolescence) and provide content that is responsive to the changing needs and capabilities of children and adolescents as they develop and establish a link to the health services [33]. Specifically, information and education about sexual activity, prevention of STIs and pregnancy, consent and bodily integrity, and promotion of healthy relationships needs to be provided before a person have sex for the first time [39–41].
Counseling and service provision for a range of modern contraceptives, with a defined minimum number and types of methods	Adolescent contraceptive use compared with adult women is characterized by shorter periods of consistent use, and greater proportions of adolescents discontinue contraceptive methods and experience contraceptive failure [44]. Many adolescents have inadequate knowledge about contraception, and they are particularly sensitive to contraceptive side effects [45,46].	CSE programmes should address gender and power dynamics including how harmful notions of masculinity and femininity affect behaviors, are perpetuated and can be transformed; rights and coercion; gender inequality in society; unequal power in intimate relationships; fostering young women's empowerment; or gender and power dynamics of condom use [43].
	Adolescents are at high risk of rapid repeat pregnancies because of misinformation or lack of awareness of fertility patterns, including return of fertility [48].	Health care providers should pay attention to and provide support for active management of side effects and continuous access to tailored information throughout the period of method use [47].
	Inadequate knowledge, skills, and judgmental attitudes of health workers (including providers who believe that adolescents should not be sexually active or that contraception may inhibit future fertility) deter service use by adolescents [50].	Contraceptive counseling and service provision should be included as an integral part of antenatal, postpartum and safe and/or postabortion care for adolescents and with timely and feasible referral mechanisms where needed [49].
	A range of health conditions often manifest for the first time in adolescence (acne, premenstrual syndrome, menstrual cramps, endometriosis symptoms, or polycystic ovarian syndrome) indicating the therapeutic use of hormonal contraceptives [54].	Adolescents should be able to access respectful and comprehensive counseling, provision of services, and/or timely and functioning referral mechanisms as required [51].
Identity formation and future thinking are two key developmental features of adolescence [55]. They include sexuality and relationships building, as well as considering aspirations and plans for the future.	Service providers should be knowledgeable about all methods, including emergency contraception and long-acting reversible contraceptives and their appropriateness for adolescents, including communicating advantages and disadvantages of methods, and the possibility of switching methods [51–53].	
	Hormonal contraception should be offered/provided to adolescents for medical reasons other than sexual activity if needed [54].	
	Contraceptive services should relate to adolescents' life goals in addition to reproductive intentions, with a defined minimum number and types of methods made available, including long-acting reversible contraception and emergency contraception [56]. Providers should allow sufficient time with their adolescent patients to address contraceptive needs using a developmentally appropriate, patient-centered approach, such as motivational interviewing [57] or aspirational counseling, meeting adolescents where they are in their life situation, and offering contraception as a means to achieving their life goals [58].	

Table 1
Continued

SRHR intervention	Related adolescent-specific attributes, needs, and barriers	Implication for service delivery
Antenatal, childbirth, and postnatal care, including emergency obstetric and newborn care	Depending on the region, although a large percentage of adolescents use some antenatal care for their first birth, they seek care later, make fewer visits during pregnancy, and receive fewer components of care than adult first-time mothers [59].	The health system should actively target and reach out to pregnant adolescents, mothers, and their families to encourage antenatal care visits and increase awareness around danger signs and when and how to seek care, including giving birth at a facility [60,61]. Health-care providers should have ongoing training and support to ensure they have the knowledge, understanding and skills to provide high-quality adolescent-responsive antenatal, intrapartum and postnatal care [60].
	Adolescent girls are likely to be influenced by social norms and practices around childbirth and may not have the autonomy to seek services according to their needs [62].	All adolescents should have information on the importance of using maternal health services through CSE, and communities need to be engaged through participatory programs [63].
	Pregnant adolescents living with HIV have lower uptake of PMTCT services than older women and their infants have poorer HIV related clinical outcomes [123].	HIV testing of pregnant adolescents during antenatal visits and differentiated services should be ensured to maintain retention [64,65].
	Perinatal depression in parenting adolescents is two to nine times higher than that in the general population, and pregnant and parenting adolescents may face several adversities such as social stigma, lack of emotional support, poor health care access, and stresses around new life adjustments [66].	Adolescent and maternal health policies should recognize mental health needs of adolescents, and health facility and/or community-based sexual and reproductive health services should link to the appropriate mental health and psychosocial support services to address related mental health conditions [67].
Safe abortion services and treatment of complications of unsafe abortion	Laws and policies in many places require girls who are pregnant to be expelled/suspended from school or does not allow them to return to school after pregnancy [68].	The legal and policy framework should allow and encourage pregnant adolescents and adolescent mothers to continue their education, and schools must be a safe space for pregnant girls and mothers [68].
	Compared with older women, adolescents are more likely to seek abortions from untrained providers or to have a self-induced abortion [30].	In countries where abortion is legal, national standards and guidelines should include both surgical and medical methods of abortion, including their application and relevance to adolescents [69].
	Adolescents know less about their rights concerning abortion and postabortion care compared with older women [30], and globally, adolescent girls are more likely than older women to delay seeking an abortion because they tend to recognize and acknowledge their pregnancy later [70].	Adolescents should be informed about their legal rights to abortion and/or post abortion care and the availability of these services at health facilities [30].
Prevention and treatment of HIV and other sexually transmitted infections	In many countries, stigma against premarital sex and pregnancy outside of marriage is pervasive, which increases adolescents' likelihood of abortion [70].	Providers should be trained to pay specific attention to confidentiality and respect in providing safe abortion care to adolescents, and be knowledgeable of when services are legal, including for adolescents [30,69].
	Adolescents have low rates of HIV and STI testing, linkage to prevention and poor treatment and adherence [71].	In generalized HIV epidemics, provider-initiated HIV testing should be offered to adolescents and youth visiting health services [72]. Health service linkages with peer and community-based ARV adherence support groups should be encouraged [73–75]. Health-care providers should be trained and supported to inform, counsel and care for adolescents according to their evolving capacities to understand the treatment and care options being offered [76].
	Adolescents living with HIV, acquired perinatally, may not be aware of their status [72].	STI prevention and management services should be provided for adolescents without mandatory parental and guardian authorization or notification [60]. Adolescents from age 12 years should be made aware of their HIV status, and additional support provided to them in disclosing their HIV status (when, how, and to whom to disclose), and confidentiality maintained [72].
	Young people who inject drugs, men who have sex with men, transgender people, sex workers, prisoners and migrants (young key populations) experience stigma, discrimination even greater than adult key populations and sometimes are subject to criminal proceedings [77].	In low and concentrated epidemics HIV testing should be available particularly in community settings to attract young key populations [72].
Prevention, detection, immediate services, and referrals for cases of sexual and gender-based violence (GBV)	Intimate partner violence (IPV) can start early, with estimates showing nearly one third of 15- to 19-year-old girls experiencing IPV [78]. IPV can increase girls' risk of unintended pregnancies and induced, unsafe abortions. In some settings, IPV also increases girls' and women's risk of acquisition of HIV and STIs.	Relevant clinical treatment for GBV should be offered to adolescent victims including emergency contraception, HIV PEP, STI treatment, and syndromic case management [79].
	Child and adolescent sexual abuse and IPV are linked to increased risk of mental health issues such as depression, posttraumatic stress disorder, and suicidal ideation and attempts [80].	Cognitive behavioral therapy with a trauma focus should be offered to children and adolescents who have been sexually abused and are experiencing adverse mental health outcomes such as symptoms of posttraumatic stress disorder and depression [79].

(continued on next page)

Table 1
Continued

SRHR intervention	Related adolescent-specific attributes, needs, and barriers	Implication for service delivery
	Female genital mutilation (FGM) [81] occurs frequently during adolescence in those countries where it is still practiced.	Deinfibulation for preventing and treating urological complications, specifically recurrent urinary tract infections and urinary retention should be offered to girls and women living with type III FGM. Cognitive behavioral therapy for girls and women living with FGM who are experiencing symptoms consistent with anxiety disorders, depression, or posttraumatic stress disorder should be considered [82]. Promoting abandonment of the practice at community-level through a combination of individual empowerment, community mobilization, and law enforcement approaches can support prevention of FGM [83].
	In some countries high numbers of adolescent girls are in formal or informal union [84].	In areas with high prevalence and/or high burden, school- and community-based interventions to inform and empower girls, in combination with interventions to influence family and community norms, in order to delay the age of marriage among girls aged <18 years should be considered [60]. Educational and employment opportunities for girls through formal and nonformal channels, in order to delay marriage until at least 18 years of age should be promoted [60]. Political leaders, planners and community leaders should be encouraged to formulate and enforce laws and policies to prohibit marriage of girls before 18 years of age [60].
Prevention, detection, and management of reproductive cancers, especially cervical cancer	A majority of adult deaths from cervical cancers can be prevented if the HPV vaccine is successfully delivered to all adolescents [85].	Awareness around HPV and the vaccine among adolescent girls, their families, teachers, and other gate keepers should be raised [86]. Legal and policy barriers to HPV vaccine provision to all adolescent girls aged 9–13 years should be removed. Girls should not be asked about their sexual activity before the vaccine is administered [87]. The roll out of the HPV vaccine to all 9- to 13-year-old girls should include mixed models comprising school and health facility –based vaccination including through integration of HPV vaccine into existing adolescent health programs, school health, and other platforms [88,89].
Information, counseling, and services for subfertility and infertility	Adolescence is an important period for developing lifelong healthy lifestyles (eg, nonsmoking, obesity prevention), which can prevent common causes of later subfertility/infertility in adulthood [90] Adolescent girls who have experienced an unsafe abortion, complicated childbirth or have been exposed to STIs may suffer from subfertility and/or secondary infertility [30]. Some childhood medical treatments and conditions such as oncologic therapy and hormonal therapy used among transgender youth, and other genetic syndromes have been shown to influence fertility. Many adolescent cancer survivors express a desire to have children in the future and have concerns about their fertility [92,93]. Providers serving adolescents who have certain illnesses or chronic conditions that threaten their fertility may not be comfortable or think it appropriate to discuss fertility preservation with them [92,93].	General sexual and reproductive health promotion (eg, safe sex to prevent STIs) and overall health promotion (eg, healthy nutrition and exercise to prevent obesity, avoiding smoking, etc) including messages around infertility and reasons for infertility [91] should be provided to all adolescents. The need for information, counseling, and services for subfertility and infertility to adolescents in service guidelines and training should be recognized especially in contexts of early marriage and childbearing, and high prevalence of STIs and unsafe abortions [91] Providers should be trained to sensitively counsel adolescents undergoing and following treatment for relevant conditions/diseases about possible effects on fertility and options (when applicable) for fertility preservation [91–93].
Information, counseling, and services for sexual health and well-being	Adolescence is a time where a person starts experiencing sexual feelings, exploring sexual behaviors, and may experience anxieties and concerns in relation to body changes, desire, pleasure, masturbation, virginity, romantic love, relationships, and marriage [94]. It is also a time with significant biological changes related to sexual maturity. Girls are less likely to negotiate terms of sex, that is, less likely to insist that their partners use condoms. Harmful masculinities may drive boys to engage in risky sexual behaviors and refuse the use contraceptives including condoms [95]. Lesbian, gay, bisexual, transgender, intersex (LGBTI) and gender nonconforming adolescents often lack adequate support systems, experience the feeling of being different and not fitting in, and are exposed to higher rates of violence, bullying and harassment, and can consequently experience mental health problems including anger, depression, sadness, stress, or worry [96,97].	Much of the strategies to ensure sexual health and well-being are captured under CSE; however, health service delivery strategies should include providing counseling to adolescents about safe, pleasurable, and consensual sex and ensuring linkages with mental health services and counseling, particularly considering adolescent-specific cognitive and behavioral transitions and establishment of identity that may result in anxiety and depression [67]. Familiarity with the psychological and medical approaches to providing care to LGBTI + and gender nonconforming adolescents including through offering gender-affirming therapies for transgender persons and options for hormonal therapy [98,122].

CSE = comprehensive sexuality education; STI = sexually transmitted infection; PMTCT = prevention of mother-to-child transmission; ARV = antiretroviral; PEP = post-exposure prophylaxis.

[28]. Access to health services is influenced by social factors at personal, family, community, and national levels [18]. To this end, adolescent SRHR interventions must recognize the risk and protective factors across these ecological levels and how they influence adolescent health outcomes [19]. Box 2 illustrates the diversity of adolescents' circumstances and experiences that influences their SRHR needs as well as their ability to access services and information. This article argues that all the interventions included in the package of essential SRHR interventions [8] should be made available to adolescents, given the diversity of adolescents' SRHR needs.

A Package of Essential SRHR Interventions for Adolescents

Ensuring that the package of essential SRHR interventions is available, accessible, and acceptable to adolescents [29] requires an approach that looks at adolescents as being biologically and socially distinct from other age groups and acknowledges that they face some specific barriers when trying to access SRHR services.

Certain aspects of SRHR are unique to adolescents (Table 1), such as menarche, initiation of sexual activity, the recommended age of HPV vaccination, and the opportunity to provide comprehensive sexuality education in school [8]. For other aspects of SRHR, the risk of adverse outcomes can be exacerbated during adolescence. This is reflected in the higher proportion of adolescent girls with an unmet need for contraception (than women aged 20–49 years married or in union), compared with older women, adolescents have a greater tendency to seek abortions from untrained providers or to self-induce; a smaller proportion of pregnant adolescent girls attending all four ANC visits compared with other women of reproductive age; adolescents being the only age group in which HIV-related deaths are not decreasing; and the greater burden of recent intimate partner violence among adolescent girls relative to their adult counterparts in several settings, including the high risk of forced sexual debut [29,30,70,78,80,99–102].

Factors increasing the risk of adverse SRHR outcomes include restrictive laws and policies, expensive or fragmented services [103], biased providers [60], and social and gender norms that do not admit adolescents' rights to autonomy over their bodies and to explore consensual, safe, and pleasurable sexual experiences [5,52,60,104,105]. Addressing these factors is critical for the success of every intervention included in the package of essential SRHR interventions. At the legal and policy level, this may require removing barriers to service provision such as compulsory parental or spousal consent for services including contraception, safe abortion and postabortion care, and HIV and STI testing and care [30,60,104,105]. At the service delivery level, critical strategies include removing or reducing costs of services for adolescents, introducing flexible opening hours adaptive to adolescents' schedules, and training and supporting providers on delivering SRHR services to adolescents in a respectful, confidential, and relevant manner [29,52,53,99,100,103,104,106,107]. At the community level, engaging parents, local leadership, and influential community members such as teachers through participatory learning and reflection, and supporting adolescent participation and leadership can generate support and increase demand for adolescent health care seeking behavior [52,60,69,104,108].

In addition to the cross-cutting strategies discussed previously, there are specific considerations in delivering each

intervention included in the essential package to adolescents. Although every intervention needs to meet the global standards for medical ethics, public health, and human rights—applicable to adolescents and adults alike—Table 1 provides a non-exhaustive list of adolescent-specific considerations for each of the interventions drawn from WHO guidelines and from published studies and evaluations in emerging areas [8].

Reaching Adolescents With a Package of Essential SRHR Interventions

Adolescents rarely seek services from a health facility, particularly not for preventive care [53,109]. To reach them with a package of essential SRHR interventions, the way the services are delivered matters therefore nearly as much as the services themselves. The Lancet Commission on Adolescent Health and Wellbeing [3] lists six main platforms, which provide opportunities to deliver health action to adolescents: health services, schools, laws and policies, communities, mHealth, and media and social marketing.

Programs that integrate SRHR across different platforms and sectors have shown promise to increase the coverage of SRHR services for adolescents [110]. Examples of integration of adolescent responsive services, coupled with adolescent-specific outreach and awareness raising interventions to address adolescent pregnancy, include countries such as Chile, the United Kingdom, Ethiopia [111], and Uruguay [112]. In these countries, programs were grounded in supportive national policies, scaled up across the health system and included different means of service delivery to cater to adolescent needs in respective contexts.

Adolescent immunization programs, such as the HPV vaccine roll out—administered in late childhood or early adolescence—also provide an opportunity to link to an integrated package of SRHR care for adolescents [87,113–115]. For example, in Togo, providers immunized girls against HPV while working hand in hand with teachers to inform girls and boys about puberty, menstrual health, and handwashing practices [116]. Although these programs focused on pregnancy prevention or vaccination rather than delivering a comprehensive approach to SRHR, the success of the delivery model points to the scaling up a package of essential SRHR interventions for adolescents in a similar manner.

Across sectors, synergies between health and education can yield particular high returns on coinvestment [117]. The growing number of adolescents enrolled in and attending school and a closing gender gap in education provides a strong case for intensifying collaboration between the health and education sector in particular. The school system is a cost-effective and equitable platform to deliver components of the package of essential SRHR interventions to adolescents [40,118]. With that said, cross-sectoral collaboration faces barriers, including differing priorities, funding streams, and timelines across sectors; and overcoming these barriers will require a shift in financial and political incentives [119].

Finally, as countries pursue UHC reforms, there are important health financing considerations for ensuring inclusion of adolescents' SRHR and adolescent health more broadly. For example, antenatal care services for adolescents are often covered, whereas contraception and safe abortion services may not be [120]. Most adolescents are particularly sensitive to costs associated with accessing SRH services [120]. Protection from

catastrophic expenditure [120] and removing cost-related barriers can increase access and utilization [121]. In practical terms, this means including adolescents in risk pools, using vouchers or other demand-side financing schemes to target subsidies to adolescents, and ensuring that adolescent-responsive service outlets are included as service providers in UHC schemes. This may involve adopting novel information and service delivery platforms such as mHealth solutions, telemedicine providers, and private sector channels, including pharmacies in strategic purchasing schemes.

Conclusion

All SRHR interventions are relevant to adolescents, and to deliver on the ICPD Programme of Action, the package of essential SRHR interventions must be made responsive to adolescents. As the health sector moves toward UHC, it must include SRHR, and it must cater to all adolescents including marginalized subgroups, such as unmarried adolescents, adolescents living with disabilities, and lesbian, gay, bisexual, transgender, intersex and other sex, and gender-diverse adolescents. This requires that the package of essential SRHR interventions is made available to all adolescents, leveraging multisectoral collaboration, strengthening accountability, and ensuring that adolescents are meaningfully engaged in the process. Successful implementation will rely on an enabling legal and policy environment and on a service delivery model that considers the determinants of adolescent SRHR, ensures integrated services, free or at very low cost, provided through a variety of platforms, respectful of the rights, and evolving capacities of adolescents.

References

- [1] United Nations. International Conference on Population and Development Programme of Action - 20th Anniversary Edition. New York: UNFPA; 2014.
- [2] WHO UNFPA and UNICEF. The reproductive health of adolescents: A strategy for action/a joint WHO/UNFPA/UNICEF statement. Geneva: World Health Organization; 1989.
- [3] Patton GC, Sawyer SM, Santelli JS, et al. Our future: A Lancet commission on adolescent health and wellbeing. *Lancet* 2016;387:2423–78.
- [4] Bundy DAP, de Silva N, Horton S, et al. Child and Adolescent Health and Development: Realizing Neglected Potential. In: Nundy D, De Silva N, Horton S, eds. DCP3 Child and Adolescent Health and Development. 3rd ed., Volume 8; 2017:1–24.
- [5] UNFPA. Unfinished Business, State of the world population. New York: UNFPA; 2019.
- [6] WHO. Health for the World's Adolescents: A second chance in the second decade. Geneva: World Health Organization; 2015.
- [7] Weiss HA, Ferrand RA. Improving adolescent health: An evidence-based call to action. *Lancet* 2019;393:1073–5.
- [8] Starrs AM, Ezeh AC, Barker G, et al. Accelerate progress—sexual and reproductive health and rights for all: Report of the Guttmacher–Lancet commission. *Lancet* 2018;391:2642–92.
- [9] WHO. Sexual health and its linkages to reproductive health: An operational approach. Geneva: World Health Organization; 2017.
- [10] UNFPA. Protecting girls' rights: The foundation for addressing motherhood in childhood. New York: UNFPA; 2013.
- [11] WHO. Global Accelerated Action for the Health of Adolescents (AA-HA!) guidance to support country implementation. Geneva: World Health Organization; 2017.
- [12] USAID Evidence in Action. Not without us! A tool for AYRH-responsive planning https://www.e2aproject.org/wp-content/uploads/TARP-1-Pager_English.pdf. Accessed November 6, 2019.
- [13] WHO. Global standards for quality health care services for adolescents. Geneva: World Health Organization; 2015.
- [14] WHO. Standards for improving the quality of care for children and young adolescents in health facilities. Geneva: World Health Organization; 2018.
- [15] WHO. Core competencies in adolescent health and development for primary care providers. Geneva: World Health Organization; 2015.
- [16] Gullo S, Galavotti C, Kuhlmann AS, et al. Effects of a social accountability approach, CARE's community score card, on reproductive health-related outcomes in Malawi: A cluster-randomized controlled evaluation. *PLoS One* 2017;12:1–20. <https://doi.org/10.1371/journal.pone.0171316>.
- [17] Braveman P, Gottlieb L. The social determinants of health: it's time to consider the causes of the causes. *Public Health Rep* 2014;129(Suppl): 19–31. <https://doi.org/10.1177/003335491412915206>.
- [18] Levesque JF, Harris MF, Russell G. Patient-centred access to health care: Conceptualising access at the interface of health systems and populations. *Int J Equity Health* 2013;12:1–40. <https://doi.org/10.1186/1475-9276-12-18>.
- [19] Viner RM, Ozer EM, Denny S, et al. Adolescence and the social determinants of health. *Lancet* 2012;379:1641–52. [https://doi.org/10.1016/S0140-6736\(12\)60149-4](https://doi.org/10.1016/S0140-6736(12)60149-4).
- [20] WHO Commission on Social Determinant of Health. Closing the gap in a generation- Health equity through action on the social determinants of health. Geneva: World Health Organization; 2008.
- [21] Dahl RE, Allen NB, Wilbrecht L, et al. Importance of investing in adolescence from a developmental science perspective. *Nature* 2018;554:441.
- [22] United Nations. Convention on the rights of the child (CRC). Off United Nations High Comm Hum Rights. 1990. <https://doi.org/10.1038/nature25770>.
- [23] Sebastian RA, Ramos MM, Stumbo S, et al. Measuring youth health engagement: Development of the youth engagement with health services survey. *J Adolesc Heal* 2014;55:334–40. <https://doi.org/10.1016/j.jadohealth.2014.02.008>.
- [24] WHO, UNFPA, UNICEF. The reproductive health of adolescents: A strategy for action. Geneva: World Health Organization; 1989.
- [25] WHO. Action for adolescent health towards a common agenda. Geneva: World Health Organization; 1997.
- [26] PMNCH, International Youth Alliance for Family Planning (IYAFP) FP 2020. Global Consensus Statement Meaningful Adolescent & Youth Engagement. Geneva: PMNCH; 2018:4–7.
- [27] USAID. A systematic review of positive youth development programs in low- and middle-income countries. Usaid 2017;1:1–132. Available at <https://www.youthpower.org/sites/default/files/YouthPower/files/resources/SystematicReview%20FINAL%209-26-17%20compress.pdf>. Accessed November 7, 2019.
- [28] Brofenbrenner U. The ecology of human development: Experiments by nature and design. Cambridge, Massachusetts: Harvard University Press; 1979.
- [29] Denno DM, Hoopes AJ, Chandra-Mouli V. Effective strategies to provide adolescent sexual and reproductive health services and to increase demand and community support. *J Adolesc Heal* 2015;56: S22–41. <https://doi.org/10.1016/j.jadohealth.2014.09.012>.
- [30] Woog V, Singh S, Browne A, et al. Adolescent women's need for and use of sexual and reproductive health services in developing countries. *New York Guttmacher Inst* 2015:1–63. Available at <https://www.guttmacher.org/report/adolescent-womens-need-and-use-sexual-and-reproductive-health-services-developing-countries>. Accessed November 7, 2019.
- [31] Ayalew M, Mengistie B, Semahegn A. Adolescent-parent communication on sexual and reproductive health issues among high school students. *Reprod Health* 2014;11:77. <https://doi.org/10.1186/1742-4755-11-77>.
- [32] Muhwezi WW, Katahoire AR, Banura C, et al. Perceptions and experiences of adolescents, parents and school administrators regarding adolescent-parent communication on sexual and reproductive health issues in urban and rural Uganda Adolescent Health. *Reprod Health* 2015;12:1–16. <https://doi.org/10.1186/s12978-015-0099-3>.
- [33] UNESCO. International technical guidance on sexuality education. Paris: UNESCO; 2018.
- [34] UNESCO Institute of Statistics. One in five children, adolescents and youth is out of school. Paris: UNESCO; 2018. Fact Sheet:13.
- [35] Ashcraft AM, Murray PJ. Talking to parents about adolescent sexuality. *Pediatr Clin North Am* 2017;64:305–20. <https://doi.org/10.1016/j.pcl.2016.11.002>.
- [36] Panchard C, Keogh SC, Stillman M, et al. Towards comprehensive sexuality education: A comparative analysis of the policy environment surrounding school-based sexuality education in Ghana, Peru, Kenya and Guatemala. *Sex Educ* 2019;19:277–96. <https://doi.org/10.1080/14681811.2018.1533460>.
- [37] Chandra-Mouli V, Patel SV. Mapping the knowledge and understanding of menarche, menstrual hygiene and menstrual health among adolescent girls in low- and middle-income countries. *Reprod Health* 2017;14:1–16. <https://doi.org/10.1186/s12978-017-0293-6>.
- [38] Wellings K, Collumbien M, Slaymaker E, et al. Sexual behaviour in context: A global perspective. *Lancet* 2006;368:1706–28. [https://doi.org/10.1016/S0140-6736\(06\)69479-8](https://doi.org/10.1016/S0140-6736(06)69479-8).
- [39] UNESCO. Review of the evidence on sexuality education report to inform the update of the UNESCO international technical guidance on sexuality education. Paris: UNESCO; 2007.

- [40] Kirby D, Laris BA, Rollieri L. Impact of sex and HIV education programs on sexual behaviors of youth in developing and developed countries. *Fam Heal Internatona*. 2005 No. 2 (Youth Research Working Paper):1–45.
- [41] Kirby D, Obasi A, Laris BA. The effectiveness of sex education and HIV education interventions in schools in developing countries. *World Health Organization - Tech Rep Ser* 2006;938:103–50.
- [42] Lundgren R, Amin A. Addressing intimate partner violence and sexual violence among adolescents: Emerging evidence of effectiveness. *J Adolesc Heal* 2015;56:S42–50. <https://doi.org/10.1016/j.jadohealth.2014.08.012>.
- [43] Haberland NA. The case for addressing gender and power in sexuality and HIV education: A comprehensive review of evaluation studies. *Int Perspect Sex Reprod Health* 2015;41:31–42. <https://doi.org/10.1363/4103115>.
- [44] Tsui AO, Trevitt JL, Blanc AK, et al. Patterns and trends in adolescents' contraceptive use and discontinuation in developing countries and comparisons. *Int Perspect Sex Reprod Health* 2009;35:63–71.
- [45] McNicholas C, Peipert JF. Long-acting reversible contraception (LARC) for adolescents. *Curr Opin Obstetr Gynecol* 2012;24:293–8. <https://doi.org/10.1097/GCO.0b013e32835686d5>.
- [46] Teal SB, Sheeder JF. IUD use in adolescent mothers: Retention, failure and reasons for discontinuation. *Contraception* 2012;85:270–4. <https://doi.org/10.1016/j.contraception.2011.07.001>.
- [47] Timmons SE, Shakibnia EB, Gold MA, et al. MyLARC: A Theory-based Interactive Smartphone App to support adolescents' use of long-acting reversible contraception. *J Pediatr Adolesc Gynecol* 2018;31:285–90. <https://doi.org/10.1016/j.jpagg.2017.11.005>.
- [48] Frarey a, Gurney EP, Sober S, et al. Postpartum contraceptive counseling for first-time adolescent mothers: A randomized controlled trial. *Arch Gynecol Obstet* 2018;361–9. <https://doi.org/10.1007/s00404-018-4969-0>.
- [49] Curtis C, Huber D, Moss-Knight T. Postabortion family planning: Addressing the cycle of repeat unintended pregnancy and abortion. *Int Fam Plan Perspect* 2010;36:44–8. <https://doi.org/10.1363/ipsrh.36.044.10>.
- [50] Ambresin AE, Bennett K, Patton GC, et al. Assessment of youth-friendly health care: A systematic review of indicators drawn from young people's perspectives. *J Adolesc Heal* 2013;52:670–81. <https://doi.org/10.1016/j.jadohealth.2012.12.014>.
- [51] Riley M, Patterson V, Lane JC, et al. The adolescent champion model: Primary care becomes adolescent-centered via targeted quality improvement. *J Pediatr* 2018;193:229–236.e1. <https://doi.org/10.1016/j.jpeds.2017.09.084>.
- [52] High-impact practices in family planning (HIPS). Adolescent-friendly contraceptive services: Mainstreaming adolescent-friendly elements into existing contraceptive services. Baltimore: John Hopkins University; 2015.
- [53] WHO. Making Health Services Adolescent Friendly. Developing National Quality Standards for Adolescent Friendly Health Services. Geneva: World Health Organization; 2012:1–44.
- [54] MacGregor KE, Khadr SN. Contraception for adolescents (*American Academy of Pediatrics*). *Arch Dis Child Educ Pract Ed* 2016;101:61–4. <https://doi.org/10.1136/archdischild-2014-307955>.
- [55] Erikson E. *Childhood and society*. Middlesex: Penguin Books; 1972.
- [56] American College of Obstetricians and Gynecologists. Committee opinion No. 735: Adolescents and long-acting reversible contraception: Implants and intrauterine devices. *Obstet Gynecol* 2018;131:e130–9.
- [57] American Academy on Pediatrics. Committee on adolescence. Contraception for adolescent policy statement. *Pediatrics* 2014;134:e1244–56.
- [58] Norton M, Chandra-Mouli V, Lane C. Interventions for preventing unintended, rapid repeat pregnancy among adolescents: A review of the evidence and lessons from high-quality evaluations. *Glob Heal Sci Pract* 2017;5:547–70. <https://doi.org/10.9745/GHSP-D-17-00131>.
- [59] Owolabi OO, Wong KLM, Dennis ML, et al. Comparing the use and content of antenatal care in adolescent and older first-time mothers in 13 countries of West Africa: A cross-sectional analysis of demographic and health surveys. *Lancet Child Adolesc Heal* 2017;1:203–12. [https://doi.org/10.1016/S2352-4642\(17\)30025-1](https://doi.org/10.1016/S2352-4642(17)30025-1).
- [60] WHO. Preventing Early pregnancy and poor reproductive outcomes. WHO Guidel. Geneva: World Health Organization; 2011:1–95. Available at https://www.who.int/immunization/hpv/target/preventing_early_pregnancy_and_poor_reproductive_outcomes_who_2006.pdf. Accessed November 7, 2019.
- [61] Salam RA, Faqqah A, Sajjad N, et al. Improving adolescent sexual and reproductive health: A systematic review of potential interventions. *J Adolesc Heal* 2016;59:S11–28. <https://doi.org/10.1016/j.jadohealth.2016.05.022>.
- [62] Morris JL, Ushwan H. Adolescent sexual and reproductive health: The global challenges. *J Gynecol Obstet* 2015;131(1):S40–2. <https://doi.org/10.1016/j.jigo.2015.02.006>.
- [63] Datiko DG, Bunte EM, Birrie GB, et al. Community participation and maternal health service utilization: Lessons from the health extension programme in rural southern Ethiopia. *J Glob Heal Rep* 2019;3:1–12.
- [64] Woldebet S, Jackson D, Lombard C, et al. Missed opportunities along the prevention of mother-to-child transmission services cascade in South Africa: Uptake, determinants, and attributable risk (the SAPMTCTE). *PLoS One* 2015;10:1–15. <https://doi.org/10.1371/journal.pone.0132425>.
- [65] UNAIDS. Start free stay free AIDS free. Geneva: UNAIDS; 2017. progress report. 2017.
- [66] Recto P, Champion JD. Psychosocial risk factors for perinatal depression among female adolescents: A systematic review. *Issues Ment Health Nurs* 2017;38:633–42. <https://doi.org/10.1080/01612840.2017.1330908>.
- [67] Hodgkinson S, Beers L, Southammakosane C, et al. Addressing the mental health needs of pregnant and parenting adolescents. *Pediatrics* 2014;133:114–22. <https://doi.org/10.1542/peds.2013-0927>.
- [68] Birchall J. Early marriage, pregnancy and girl child school dropout. *Knowledge, Evid Learn Dev* 2018;2:1–15.
- [69] WHO. Safe abortion :technical and policy guidance for health systems. 2nd ed. Geneva: World Health Organization; 2012.
- [70] Guttmacher Institute. Abortion worldwide 2017: Uneven progress and unequal access. New York: Guttmacher Institute; 2017. Available at <https://www.guttmacher.org/report/abortion-worldwide-2017>. Accessed November 7, 2019.
- [71] Wong VJ, Murray KR, Phelps BR, et al. Adolescents, young people, and the 90-90-90 goals: A call to improve HIV testing and linkage to treatment. *Aids* 2017;31:S191–4. <https://doi.org/10.1097/QAD.0000000000001539>.
- [72] WHO. HIV and adolescents: Guidance for HIV testing and counselling and care for adolescents living with HIV: Recommendations for a public health approach and considerations for policy-makers and managers, World Health. Geneva: World Health Organization; 2013:171.
- [73] Ridgeway K, Dulli LS, Murray KR, et al. Interventions to improve antiretroviral therapy adherence among adolescents in low- and middle-income countries: A systematic review of the literature. *PLoS One* 2018;13:e0189770. <https://doi.org/10.1371/journal.pone.0189770>.
- [74] Willis N, Napei T, Armstrong A, et al. Zvandiri—Bringing a differentiated service delivery program to scale for children, adolescents, and young people in Zimbabwe. *JAIDS J Acquir Immune Defic Syndr* 2018;78:S115–23. <https://doi.org/10.1097/QAI.0000000000001737>.
- [75] WHO. What's new in adolescent treatment and care fact sheet: HIV treatment and care. Geneva: World Health Organization; 2016.
- [76] WHO. Global health sector strategy on sexually transmitted infections 2016–2021, 1. Geneva: World Health Organization; 2016:1–63.
- [77] Interagency Working Group on Key Populations. HIV and young key populations: A technical brief series. Geneva: World Health Organization; 2016.
- [78] WHO London School of Hygiene and Tropical Medicine South African Medical Research Council. Global and regional estimates of violence against women: Prevalence and health effects of intimate partner violence and non-partner sexual violence. Geneva: World Health Organization; 2013.
- [79] WHO. Responding to children and adolescents who have been sexually abused. Geneva: World Health Organization; 2017.
- [80] WHO. WHO recommendations on adolescent sexual and reproductive health and rights. Geneva: World Health Organization; 2018.
- [81] UNICEF. Female genital mutilation. Available at: <https://data.unicef.org/topic/child-protection/female-genital-mutilation/>. Accessed September 15, 2019.
- [82] WHO. WHO guidelines on the management of health complications from female genital mutilation. WHO Guidel Manag Heal Complicat Female Genit Mutilation 2016;47.
- [83] UNFPA, UNICEF. How to transform a social norm. New York: UNFPA, UNICEF; 2018.
- [84] UNFPA. Marrying too young. New York: UNFPA; 2012.
- [85] Wardak S. Human Papillomavirus (HPV) and cervical cancer. *Med Dosw Mikrobiol* 2016;68:73–84.
- [86] Snidal S. Lessons learned from Uganda: Delivering HPV vaccines to hard-to-reach girls | Clinton Health Access Initiative. *Clint Heal Access Initiat* 2018. Available at <https://clintonhealthaccess.org/lessons-learned-uganda-delivering-hpv-vaccines-hard-reach-girls/>. Accessed November 19, 2019.
- [87] WHO. Comprehensive cervical cancer control. Geneva: World Health Organization; 2014:366–78.
- [88] Ladner J, Besson MH, Hampshire R, et al. Assessment of eight HPV vaccination programs implemented in lowest income countries. *BMC Public Health* 2012;12:1. <https://doi.org/10.1186/1471-2458-12-370>.
- [89] Pollack AE, Balkin M, Edouard L, et al. Ensuring access to HPV vaccines through integrated services: A reproductive health perspective. *Bull World Health Organ* 2007;85:57–63.
- [90] Rossi BV, Abusief M, Missmer SA. Modifiable risk factors and infertility: What are the connections? *Am J Lifestyle Med* 2016;10:220–31. <https://doi.org/10.1177/1559827614558020>.

- [91] Wimberly YH, Kahn JA, Kollar LM, et al. Adolescent beliefs about infertility. *Contraception* 2003;68:385–91. <https://doi.org/10.1016/j.contraception.2003.07.006>.
- [92] Nahata L, Quinn GP. Fertility preservation in young males at risk for infertility: What every pediatric provider should know. *J Adolesc Heal* 2017;60:237–8. <https://doi.org/10.1016/j.jadohealth.2016.12.015>.
- [93] Benedict C, Shuk E, Ford JS. Fertility issues in adolescent and young adult cancer survivors. *J Adolesc Young Adult Oncol* 2016;5:48–57. <https://doi.org/10.1089/jayao.2015.0024>.
- [94] Angela Oswalt, MSW, edited by C. E. Zupanick PD. *The Development of Adolescent Sexuality*. 2005.
- [95] Philpott A, Knerr W, Maher D. Promoting protection and pleasure: Amplifying the effectiveness of barriers against sexually transmitted infections and pregnancy. *Lancet* 2006;368:2028–31. [https://doi.org/10.1016/S0140-6736\(06\)69810-3](https://doi.org/10.1016/S0140-6736(06)69810-3).
- [96] Baltag V, Sawyer SM. Quality health care for adolescents. *Int Handb Adolesc Heal Dev Public Heal Response* 2016:309–24.
- [97] Hillier L, Jones T, Monagle M, et al. Writing themselves in 3: The third national study on the sexual health and wellbeing of same sex attracted and gender questioning young people, Monogr Ser. Bundoora: Australian Research Centre in Sex, Health and Society, La Trobe University; 2010:1–134.
- [98] World Health Organization. Consolidated guidelines on HIV prevention, diagnosis, treatment and care for key populations. *WHO Guidel* 2014:184.
- [99] Zuurmond MA, Geary RS, Ross DA. The effectiveness of youth centers in increasing use of sexual and reproductive health services: A systematic review. *Studies in family planning, Stud Fam Plann* 2012;43:239–54. <https://doi.org/10.1111/j.1728-4465.2012.00324.x>.
- [100] Hainsworth G, Engel D, Simon C, et al. Scale-up of adolescent contraceptive Services : Lessons from a 5-country comparative analysis. *J Acquir Immune Defic Syndr* 2014;66:200–8. <https://doi.org/10.1097/QAI.0000000000000180>.
- [101] Shah IH, Ahman E. Unsafe abortion differentials in 2008 by age and developing country region : High burden among young women unsafe abortion differentials in 2008 by age and developing. *RHM* 2012;20:169–73.
- [102] Decker MR, Latimore AD, Yasutake S, et al. Gender-based violence against adolescent and young adult women in low- and middle-income countries. *J Adolesc Heal* 2015;56:188–96.
- [103] Francis JKR, Gold MA. Long-Acting reversible contraception for adolescents: A review. *JAMA Pediatr* 2017;171:694–701.
- [104] WHO. Selected practice recommendations for contraceptive use. 3rd ed. Geneva: World Health Organization; 2016:1–66.
- [105] WHO. WHO consolidated guideline on self-care interventions for health: Sexual and reproductive health and rights. Geneva: World Health Organization; 2019.
- [106] Felitti VJ, Anda RF, Nordenberg D, et al. Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The adverse childhood experiences (ACE) study. *Am J Prev Med* 2019;56:774–86.
- [107] WHO. Sexually transmitted infections: Implementing the global STI strategy. Geneva: World Health Organization; 2017.
- [108] USAID- Office of Population and Reproductive Health Bureau for Global Health. Essential considerations for engaging men and boys for improved family planning outcomes. 2018.
- [109] Irwin CE, Adams SH, Park MJ, et al. Preventive care for adolescents: Few get visits and fewer get services. *Pediatrics* 2009;123.
- [110] Rasanathan K, Damji N, Atsbeha T, et al. Ensuring multisectoral action on the determinants of reproductive, maternal, newborn, child, and adolescent health in the post-2015 era. *BMJ* 2015;351:h4213.
- [111] Chandra Mouli V, Plesons M, Hadley A, et al. Lessons learned from national government- led efforts to reduce adolescent pregnancy. *Early Childhood Matters*; 2017. 50–6.
- [112] Chandra-Mouli V, Ferguson BJ, Plesons M, et al. The political, research, programmatic and social responses to adolescent sexual and reproductive health and rights in the 25 years since the International Conference on Population and Development. *J Adolesc Health* 2019;65:S16–40.
- [113] Watson-Jones D, Lees S, Mwangi J, et al. Feasibility and acceptability of delivering adolescent health interventions alongside HPV vaccination in Tanzania. *Health Policy Plan* 2016;31:691–9.
- [114] Macphail C, Venables E, Rees H, et al. Using HPV vaccination for promotion of an adolescent package of care: Opportunity and perspectives. *BMC Public Health* 2013;13:1.
- [115] WHO. Options for linking health interventions for adolescents. Geneva: World Health Organization; 2008.
- [116] GAVI. Gavi Progress Report 2016. Geneva: GAVI; 2017:47.
- [117] Bundy DAP, de Silva N, Horton S, et al. Investment in child and adolescent health and development: Key messages from disease control priorities, 3rd edition. *Lancet* 2018;391:687–99.
- [118] Shackleton N, Jamal F, Viner RM, et al. School-based interventions going beyond health education to promote adolescent health: Systematic review of reviews. *J Adolesc Heal* 2016;58:382–96.
- [119] Arcaya M, de Briggs XS. Despite obstacles, considerable potential exists for more robust federal policy on community development and health. *Health Aff* 2011;30:2064–71.
- [120] Waddington C, Sambo C. Financing health care for adolescents: A necessary part of universal health coverage. *Bull World Health Organ* 2015;93:57–9.
- [121] Meuwissen LE, Gorter AC, Knottnerus AJA. Impact of accessible sexual and reproductive health care on poor and underserved adolescents in Managua, Nicaragua: A quasi-experimental intervention study. *J Adolesc Heal* 2006;38:56.e1-56.e9.
- [122] Vance SR, Ehrensaft D, Rosenthal SM. Psychological and medical care of gender nonconforming youth. *Pediatrics* 2014;134(6):1184–92.
- [123] Horwood C, Butler LM, Haskins L, et al. HIV-Infected Adolescent Mothers and Their Infants: Low Coverage of HIV Services and High Risk of HIV Transmission in KwaZulu-Natal, South Africa. *PLoS One* 2013. <https://doi.org/10.1371/journal.pone.0074568>.